

**Running Head:**

Interactions between freezing tolerance and flowering

**Corresponding author:**

Jorge Dubcovsky

One Shields Avenue

Department of Plant Sciences

University of California

Davis, CA, 95616, USA

Phone: (530) 752-5159

FAX: (530) 752-4361

Email: [jdubcovsky@ucdavis.edu](mailto:jdubcovsky@ucdavis.edu)

**Journal Research Area:**

Development and Hormone Action – Associate Editor Richard Amasino

## **Regulation of freezing tolerance and flowering in temperate cereals: the *VRN-1* connection**

Taniya Dhillon<sup>1\*</sup>, Stephen P. Pearce<sup>2\*</sup>, Eric J. Stockinger<sup>1\*</sup>, Assaf Distelfeld<sup>2</sup>, Chengxia Li<sup>2</sup>, Andrea K. Knox<sup>1\*\*</sup>, Ildikó Vashegyi<sup>3</sup>, Attila Vágújfalvi<sup>3</sup>, Gabor Galiba<sup>3,4</sup>, Jorge Dubcovsky<sup>2§</sup>

<sup>1</sup>Department of Horticulture and Crop Science, The Ohio State University/OARDC, Wooster, OH, 44691, USA

<sup>2</sup>Department of Plant Sciences, University of California, Davis, CA, 95616, USA

<sup>3</sup>Agricultural Research Institute of the Hungarian Academy of Sciences, 2462, Martonvásár, Hungary

<sup>4</sup>Faculty of Information Technology, Research Institute of Chemical and Process Engineering, University of Pannonia, 8200, Veszprém, Hungary

\*These authors contributed equally to this work

\*\*Present address: Dept. of Biology, Williams College, Williamstown, MA, 01267, USA

§To whom correspondence should be addressed

Email: [jdubcovsky@ucdavis.edu](mailto:jdubcovsky@ucdavis.edu)

Phone: 1-530-752-5159

Fax: 1-530-752-4361

**Footnotes:**

Work in Dr. Stockinger's lab was supported in part by grants from the NSF Plant Genome Program (DBI 0110124) and the Ohio Plant Biotechnology Consortium (Proposal Number 2007-020). Work in Dr. Dubcovsky's lab was supported by the National Research Initiative Competitive Grant no. 2008-35100-04562 from the USDA National Institute of Food and Agriculture. Work in Dr. Galiba's lab was supported by the Hungarian Research Fund 'OTKA' No. K68894 and K75528.

**Corresponding author:**

Jorge Dubcovsky

FAX: (530) 752-4361

Email: [jdubcovsky@ucdavis.edu](mailto:jdubcovsky@ucdavis.edu)

## ABSTRACT

In winter wheat and barley varieties, long exposures to non-freezing cold temperatures accelerate flowering time (vernalization) and improve freezing tolerance (cold acclimation). However, when plants initiate their reproductive development, freezing tolerance decreases suggesting a connection between the two processes. To better understand this connection we used two diploid wheat mutants “*maintained vegetative phase*” (*mvp*) that carry deletions encompassing *VRN-1*, the major vernalization gene in temperate cereals. Homozygous *mvp/mvp* plants never flower, whereas plants carrying at least one functional *VRN-1* copy (*Mvp/-*) exhibit normal flowering and high transcript levels of *VRN-1* under long days. The *Mvp/-* plants showed reduced freezing tolerance and reduced transcript levels of several cold induced *CBF* transcription factors and *COLD REGULATED* genes (*COR*) relative to the *mvp/mvp* plants. Diploid wheat accessions with mutations in the *VRN-1* promoter resulting in high transcript levels under both long and short days, showed a significant down-regulation of *COR14b* under long but not under short days. Taken together these studies suggest that *VRN-1* is required for the initiation of the regulatory cascade that down-regulates the cold acclimation pathway, but that additional genes regulated by long days are required for the down-regulation of the *COR* genes. In addition, our results show that allelic variation in *VRN-1* is sufficient to determine differences in freezing tolerance suggesting that QTL for freezing tolerance previously mapped on this chromosome region are likely a pleiotropic effect of *VRN-1* rather than the effect of a separate closely linked locus (*FROST RESISTANCE-1*) as proposed in early freezing tolerance studies.

## INTRODUCTION

Exposure to low non-freezing temperatures, a process known as cold acclimation, increases a plant's freezing tolerance (Thomashow, 1990; Thomashow, 1999). Freezing tolerant plants that have not been cold acclimated are generally killed at approximately -3 to -5 °C while cold acclimated plants can survive much colder freezing temperatures. In addition, increasing the length of the cold acclimation period, up to a point, can also increase freezing tolerance. These two observations suggest that cold acclimation is an active process.

Freezing tolerance is essential for fall-planted temperate cereals (wheat, barley, and rye) to survive freezing temperatures during the winter. In contrast, spring-sown genotypes do not require high levels of freezing tolerance since they are not exposed to the freezing temperatures of winter. One feature that distinguishes winter and spring genotypes is the requirement of the former for a long period (several weeks) at cold temperature to accelerate the transition from the vegetative growth phase to the reproductive growth phase, a process called vernalization. Spring genotypes do not have a vernalization requirement and flower in the absence of the extended low temperature treatment (reviewed in Trevaskis et al., 2007; Distelfeld et al., 2009).

The requirement for exposures to non-freezing cold temperatures is common to both cold-acclimation and vernalization, suggesting a potential connection between these two processes. Winter genotypes maintained under continuous cold, after an initial increase in freezing tolerance, exhibit a progressive decrease in their cold acclimation ability (Fowler et al., 1996a; Fowler et al., 1996b; Fowler and Limin, 2004). This progressive decrease inversely parallels the fulfillment of the vernalization requirement. A clear decrease in freezing tolerance occurs after the shoot apical meristem (SAM) advances to the double ridge stage (Fowler et al., 1996a, 1999; Limin and Fowler, 2006). These studies suggest that a regulatory component of freezing tolerance is linked to a developmental shift between the vegetative and reproductive stages. Limin and Fowler (2006) suggested that the main vernalization gene, *VRN-1* that is induced during vernalization, plays an important role in the decrease of the ability to cold acclimate with development.

Early genetic studies also revealed a correlation between growth habit and freezing tolerance; wheat genotypes having a spring growth habit were less freezing tolerant than genotypes having a winter growth habit (Hayes and Aamodt, 1927). Subsequent studies carried out using wheat chromosome substitution lines revealed that homologous group 5 chromosomes, where *VRN-1* is located, have the largest effect (Roberts, 1986). The first major locus affecting freezing tolerance and winter hardiness on homoeologous group 5 was designated *FROST RESISTANCE-1 (FR-1)* (Sutka and Snape, 1989). However, since *FR-1* co-segregates with *VRN-1* in most genetic studies, it is still not clear if *FR-1* is an independent gene or just a pleiotropic effect of *VRN-1* (Brule-Babel and Fowler, 1988; Sutka and Snape, 1989; Roberts, 1990; Hayes et al., 1993; Francia et al., 2004; Galiba et al., 2009).

More recently, a second locus associated with natural variation in freezing tolerance in wheat and barley was mapped on the long arm of homoeologous group 5. This locus, designated *FROST RESISTANCE-2 (FR-2)*, is approximately 30 cM proximal to *VRN-1* and includes a cluster of 11 (or more) *C-repeat Binding Factor (CBF)* genes (Vágújfalvi et al., 2003; Francia et al., 2004; Miller et al., 2006; Skinner et al., 2006; Francia et al., 2007; Knox et al., 2010). The *FR-2 CBF* gene cluster has surfaced as a major QTL affecting freezing tolerance in a number of wheat and barley mapping populations (Vágújfalvi et al., 2003; Francia et al., 2004; Båga et al., 2007; Francia et al., 2007).

The role of the *CBF* genes in freezing tolerance has been studied in detail in *Arabidopsis thaliana*. The CBFs are transcriptional activators that promote expression of genes whose upstream regulatory sequences harbor the CRT/DRE low temperature cis-acting DNA regulatory element (Stockinger et al., 1997). Approximately 20% of the *Arabidopsis* genes whose expression is altered during cold acclimation are directly or indirectly controlled by the CBF transcription factors (Vogel et al., 2005). Direct targets of the CBFs in *Arabidopsis* include the robustly-induced *Cold Regulated (COR)* genes (Jaglo-Ottosen et al., 1998). Similar candidate *CBF* target genes in the cereals, which also harbor CRT/DRE motifs in their upstream regulatory region, include *COR14b*, *DHN5*, and *DHN8* (Dal Bosco et al., 2003; Choi et al., 1999). Many of these *COR* genes are induced to higher levels in genotypes exhibiting greater freezing tolerance than in those having lesser freezing tolerance (Houde et al., 1992; Danyluk et al., 1994; Crosatti et al.,

1996; Fowler et al., 1996b; Limin et al., 1997; Danyluk et al., 1998; Grossi et al., 1998; NDong et al., 2002). The use of *COR14b* as an expression QTL to map loci affecting *COR* expression levels revealed two major loci; one of which is coincident with *VRN-1* and the second one with *FR-2* (Vágújfalvi et al., 2000; Francia et al., 2004).

Notably, genotypes carrying the *vrn-1* allele for winter growth habit express certain *CBF* genes at higher levels than genotypes carrying the *Vrn-1* allele for spring growth habit (Stockinger et al., 2007). Moreover, once the winter genotypes carrying the *vrn-1* allele are vernalized, *CBF* transcript levels are dampened relative to levels detected in non-vernalized plants (Stockinger et al., 2007). This suggests that *VRN-1* somehow acts to repress expression of the *CBFs* at *FR-2*, and in turn decrease freezing tolerance.

The molecular isolation of *VRN-1* revealed that this gene encodes a MADS-box protein similar to the Arabidopsis meristem identity gene *APETALA1 (API)* (Danyluk et al., 2003; Trevaskis et al., 2003; Yan et al., 2003). The characterization of *VRN-1* alleles associated with winter and spring genotypes showed that the primary differences were insertions and deletions in regulatory regions located in the promoter and first intron (Yan et al., 2004; Fu et al., 2005; Pidal et al., 2009). Deletions in the *VRN-1* promoter affecting a small region tentatively designated the ‘VRN-box’ (Pidal et al., 2009) or large deletions / insertions in the *VRN-1* first intron are both associated with spring growth habit (Pidal et al., 2009; Fu et al., 2005). Genotypes with a winter growth habit (*vrn-1* allele) show very low *VRN-1* transcript levels until plants are vernalized. In contrast, spring genotypes (*Vrn-1* allele) constitutively express *VRN-1* to high levels. Flowering is initiated once *VRN-1* transcripts reach a critical threshold level (Loukoianov et al., 2005).

In addition to vernalization, photoperiod also plays a role in *VRN-1* regulation. In photoperiod sensitive genetic backgrounds long day photoperiods enhance *VRN-1* transcript accumulation while short day photoperiods delay transcript accumulation. In both wheat and barley the delay in the transition to floral initiation in plants grown under SD photoperiods is associated with increased freezing tolerance (Limin and Fowler, 2006; Fowler et al., 2001). One of the cis-elements responsible for the lack of *VRN-1* expression under short days in diploid wheat is thought to reside within the *VRN-1* promoter CArG motif, a binding site for MADS-box transcription factors located

downstream of the VRN-box (Dubcovsky et al., 2006; Pidal et al., 2009). Under short days, diploid wheat (*Triticum monococcum* L.) plants carrying deletions in this CArG motif (*Vrn-1f* and *Vrn-1g* alleles) show accumulation of *VRN-1* transcripts and a slow transition of the shoot apical meristem to the reproductive stage, whereas plants with intact CArG motifs (e.g. *Vrn-1h* and *vrn-1*) show no *VRN-1* expression in SD and remain in the vegetative phase (Dubcovsky et al., 2006). Under long days, all accessions with *VRN-1* alleles for spring growth habit show accumulation of *VRN-1* transcripts and a rapid initiation of the transition to the reproductive stage (Dubcovsky et al., 2006).

*Triticum monococcum* mutants with deletions of the *VRN-1* gene fail to flower indicating that this gene is indispensable for the transition to the reproductive phase (Shitsukawa et al., 2007a). Two independently induced nitrogen-ion-beam mutants, designated *maintained vegetative phase 1* and *2* (*mvp-1* and *mvp-2*) were generated in different *T. monococcum* genetic backgrounds. The deletions in these two mutants encompass the complete *VRN-1* gene and several closely linked genes (Distelfeld and Dubcovsky, 2010).

To investigate the role of *VRN-1* in freezing tolerance we made use of the *mvp* mutants and of natural *T. monococcum* accessions that differ in their ability to express *VRN-1* under short days. We found that freezing tolerance and transcript levels of several *CBF* and *COR* genes were higher in the *mvp* mutants relative to the plants carrying at least one functional *VRN-1* copy. However, the expression of *VRN-1* under short days was not as effective as under long days to down-regulate *COR14b* gene transcription. Taken together these results suggest that *VRN-1* transcription is necessary but not sufficient to down-regulate the *COR* genes.

## RESULTS

### Effect of the *mvp* mutations on freezing tolerance

The homozygous *mvp-2* mutants (*mvp-2/mvp-2*) and the plants carrying at least one functional copy of *VRN-1* (*Mvp-2/-*) were identified using a dominant molecular marker

for *VRN-1* as described in the SOM (Fig. S1). Just before the freezing experiments, there were clear differences in apical development between plants from each group. The apices from the *mvp-2/mvp-2* mutant were at the vegetative stage whereas those from the *Mvp-2/-* plants were already at the double ridge stage (Fig. S2).

Significant differences in survival rates were detected between the *mvp-2/mvp-2* and *Mvp-2/-* plants in controlled freezing experiments. Differences between the two genotypic classes were detected at both -9 °C and -11 °C freezing temperatures (Table 1). In the group frozen to -9 °C none of the *Mvp-2/-* plants survived, whereas 87% of the *mvp-2/mvp-2* mutants survived (Table 1). In the group frozen to -11 °C none of the *Mvp-2/-* plants survived, whereas approximately half (46%) of the *mvp-2/mvp-2* mutants survived (Table 1). In a second freezing experiment performed under slightly different acclimation and freezing conditions (SOM) approximately 70% of the *mvp-2/mvp-2* mutants survived -12 °C freezing temperatures whereas only 40% of the *Mvp-2/-* plants survived the same treatment ( $P=0.009$ , SOM, Fig. S3). In this second experiment, all plants from both genotypic classes were killed at -13 °C (Fig. S2, SOM).

To evaluate the effect of freezing on the functionality of the Photosystem II (PSII), the maximum quantum yield of the PSII photochemistry was measured by the ratio of variable ( $F_v$ ) to maximal ( $F_m$ ) fluorescence in a dark-adapted state,  $F_v/F_m$  (Butler and Kitajima 1975).  $F_v/F_m$  ratios taken 2-4 hours after returning the plants to 20 °C paralleled the survival results (high  $F_v/F_m$  values indicate low freezing damage). Homozygous *mvp-2/mvp-2* mutant plants showed significantly higher  $F_v/F_m$  values than the *Mvp-2/-* plants (Table 1,  $P<0.0001$ ), in agreement with the greater freezing tolerance of the *mvp-2/mvp-2* plants.

Along with the  $F_v/F_m$  measurements, samples of the crown and adjoining tissues were collected to measure relative conductivity. This measurement estimates the cellular electrolytes leached from the freeze-damaged tissue as a proportion of the total cellular electrolytes and is based on the principle that the greater the damage to cells from freezing injury, the greater the exosmosis of cellular electrolytes into a water solvent (Dexter, 1956). The *Mvp-2/-* plants showed higher relative conductivity values than those from the homozygous *mvp-2/mvp-2* mutants, both at -9 °C (19% increase) and -11 °C

(45% increase). However, the differences between genotypes were only marginally significant (Table 1,  $P=0.05$ ), likely due to the limited number of mutant plants sacrificed for relative conductivity measures in this first experiment (three *mvp-2/mvp-2* and six *Mvp-2/-* plants per temperature). A second experiment using leaves from 9 plants per genotype-temperature combination confirmed the higher relative conductivity of the *Mvp-2/-* plants compared with the *mvp-2/mvp-2* (76% average increase over the three temperatures,  $P=0.003$ , SOM Figure S3).

Taken together these three sets of data indicate that the presence of the *VRN-1* gene in the *Mvp-2/-* lines is associated with a decrease in freezing tolerance relative to the *mvp* mutants.

### **Effect of the *mvp* mutations on *CBF* transcript levels**

Quantitative RT-PCR was used to compare transcript levels of 11 *CBF* genes in 4 week-old *Mvp-2/-* and *mvp-2/mvp-2* plants both before (20 °C) and after 8 h of cold treatment at 4 °C (Fig. 1). With the exception of *CBF2*, which showed very low levels of expression both at 20 °C and 4 °C, the other 10 *CBF* genes showed very low transcript levels at 20 °C and were significantly up-regulated after 8 h of cold treatment ( $P<0.01$ ). Five *CBF* genes (*CBF2*, *CBF4*, *CBF9*, *CBF12*, *CBF17*) showed significantly higher expression levels ( $P<0.01$ ) in the *mvp-2/mvp-2* homozygous mutant plants than in the *Mvp-2/-* plants (Fig. 1). The same difference was marginally significant for *CBF14* ( $P=0.03$ ) and not significant for the other *CBF* genes.

A second experiment was carried out using 8-week old *Mvp-1/-* and *mvp-1/mvp-1* plants and, as before, measurements of *CBF* transcript levels after 8 h at 4 °C (SOM, Fig. S4). However, in this second experiment no samples were collected at 20 °C because of the negligible *CBF* transcript levels observed in the first experiment at this temperature. In the *Mvp-1/-* plants the shoot apical meristems were between the double ridge and terminal spikelet stages, whereas in the *mvp-1/mvp-1* mutants the shoot apical meristems were in the vegetative stage. As in the 4-week old *Mvp-2* plants used in the previous experiment, the older 8-week *Mvp-1* plants showed significantly lower transcript levels of *CBF2*, *CBF4*, *CBF9*, *CBF12*, and *CBF17* in the *Mvp-1/-* plants relative to the

homozygous *mvp-1/mvp-1* mutants after the cold treatment. Whereas *CBF14* transcript level differences between *Mvp-2/-* and *mvp-2/mvp-2* genotypes were marginally significant in the first experiment, in this second experiment the differences in *CBF14* transcript levels between *Mvp-1/-* and *mvp-1/mvp-1* were not significant. Curiously, transcript levels of *CBF12* and *CBF16* were much lower relative to the other genes in the 8-week old *Mvp-1* plants relative to the 4-week old *Mvp-2* plants.

Quantitative PCR measurements of *VRN-1* in both experiments revealed high levels of *VRN-1* transcripts in the *Mvp-1/-* and *Mvp-2/-* plants, which were even higher than those of the highly expressed *TEF1* endogenous control gene in both cases. As expected, no *VRN-1* transcripts were detected in the homozygous *mvp/mvp* mutants. Taken together, these results suggest that the presence of *VRN-1* (or genes regulated by *VRN-1*) modulates the response of several *CBF* genes to cold.

### **Effect of the *mvp* mutations on *COR* gene transcript levels**

Eight hours after transferring 8-week old plants from 20 °C to 4 °C, *COR14b* transcripts were two-fold higher ( $P < 0.05$ ) in the homozygous *mvp-1/mvp-1* plants than in the *Mvp-1/-* plants (Fig. S4). Two additional experiments, in which plants were exposed to longer periods of cold temperatures, were performed to further characterize the differences in *COR14b* transcript levels between mutants and non-mutants.

In the first experiment, four-week old *mvp-2/mvp-2* and *Mvp-2/-* plants were transferred from room temperature to 4 °C and kept at that temperature for 12 days. Leaf samples for RNA analysis were collected the day prior to the cold treatment and 8 h, 32 h, 4 days, and 12 days after transferring the plants to 4 °C. All samples were collected at 2 pm to avoid differences that might be caused by circadian effects. In this qRT-PCR experiment, transcript levels from both genotypes peaked at 32 h and then decayed slowly during the next 11 days. At each of the last three sampling points the *COR14b* transcript levels were significantly higher ( $P < 0.001$ ) in the *mvp-2/mvp-2* homozygous plants than in *Mvp-2/-* plants (Fig. 2).

In the second experiment, steady state transcript levels of *COR14b* and two additional *COR* genes (*DHN5* and *DHN8*) were evaluated by RNA blot analysis using a more gradual decrease in temperatures and longer time exposures to the inductive temperatures (19 d at 10 °C followed by 12 d at 4 °C, all under long days) (Fig. 3). At the beginning of the cold induction the *Mvp-2/-* plants were already induced to flower and showed high levels of *VRN-1* transcripts whereas the *mvp-2/mvp-2* plants were in the vegetative stage and, as expected from the homozygous deletion, showed no *VRN-1* transcripts (Fig. 3).

At all time points after the cold induction, the transcript levels of *COR14b* and *DHN5* were higher in the *mvp-2/mvp-2* plants (no *VRN-1* transcripts) than in the *Mvp-2/-* (high *VRN-1* transcripts). The *DHN8* gene did not show this alternate pattern between genotypes (Fig. 3). In *mvp-2/mvp-2* homozygous mutants *COR14b* and *DHN5* levels remained high throughout the sampling time course (Fig. 3).

Taken together, the expression data from these experiments showed that in plants with high *VRN-1* transcript levels several *CBF* and *COR* genes are down-regulated. To test if the down-regulation of these genes was a direct effect of the increase in *VRN-1* transcript levels or a result of the transition to the reproductive phase, the relationship between *VRN-1* and *COR14b* transcript levels was studied in *T. monococcum* lines that, under short days, differ in the expression of *VRN-1* but show similar delays in the progression to the reproductive phase.

### **Effect of *VRN-1* transcription on *COR14b* transcript levels under short days**

*Triticum monococcum* lines carrying a “wild-type” *vrn-A1* allele and recessive *vrn-A2* alleles have a spring growth habit and show no expression of *VRN-1* under short days (Dubcovsky et al., 2006). However, *T. monococcum* lines carrying a 1-bp deletion in the *VRN-1* promoter CArG box and an insertion in intron one (*Vrn-1f* allele), or a 34-bp deletion encompassing the complete CArG box (*Vrn-1g* allele), show high *VRN-1* transcript levels under short days (Dubcovsky et al., 2006). These plants show a transition of the shoot apical meristem to the double ridge stage under short days, but further development of the spike is delayed until the plants are transferred to long days (Dubcovsky et al., 2006).

Expression profiling of three independent accessions of *T. monococcum* lines carrying each of the three genotypes confirmed previously published results (Dubcovsky et al., 2006). Under short days, lines having the *Vrn-1f* or *Vrn-1g* alleles showed high *VRN-1* transcript levels whereas those having the wild-type *vrn-1* allele showed no *VRN-1* transcripts (Fig. 4A). As expected, transcript levels of *VRN-1* were higher at six weeks than at two weeks, and *COR14b* transcripts were high after cold temperatures and nearly absent in plants maintained at warm temperature (Fig. 4A).

In the six-week old *Vrn-1f* and *Vrn-1g* lines grown under short days, *COR14b* was highly responsive to cold temperatures, despite relatively high *VRN-1* transcript levels in these lines (Fig. 4A, arrows). In contrast, when six-week old plants of these same genotypes were grown under long day conditions, the same cold treatment failed to induce *COR14b* to high levels (Fig. 4A). This result indicates that under short days the expression of *VRN-1* was not sufficient to down-regulate *COR14b*.

Analyses of the two-week old plants (Fig. 4A) further confirmed the inverse correlation between *VRN-1* and *COR14b*. Under short days, the cold treatments resulted in a strong up-regulation of *COR14b* (low *VRN-1* transcript levels in all genotypes), but under long days the down-regulation of *COR14b* was not as strong as in the 6-week old plants, likely because the *VRN-1* transcript levels in two-week-old plants was not as high as in the 6-week old plants (Fig. 4A). To confirm these results this experiment was repeated using qRT-PCR and the same genotypes at the same developmental stage (Fig. 4B). In plants grown under short days and exposed for 32 h to 4°C, *COR14b* transcript levels showed no-significant differences ( $P= 0.95$ ) between wild-type and *Vrn-1* mutant lines (*Vrn-1f* and *Vrn-1g*), despite significant differences in *VRN-1* transcript levels. In contrast, highly significant differences in *COR14b* transcript levels ( $P= 0.006$ ) were detected under long days, in which *COR14b* transcript levels in the *Vrn-1f* and *Vrn-1g* mutant lines were lower than in the wild-type (Fig. 4B).

An additional experiment was carried out to compare *COR14b* expression levels in *T. monococcum* lines carrying the *Vrn-1f* and *Vrn-1g* alleles (early spring growth habit) to those in a *T. monococcum* accession carrying the *Vrn-1h* allele (late spring growth habit). The *Vrn-1h* allele has an almost identical repetitive element insertion in the first intron as

*Vrn-1f*, but lacks the CArG box mutation (Dubcovsky et al. 2006). As previously reported the line with the *Vrn-1h* allele showed low *VRN-1* transcripts under short days and maintained low *VRN-1* transcript levels even when plants were grown under long days for six weeks. The results presented in Fig. 4C confirmed that high *VRN-1* transcript levels under short days (*Vrn-1f* and *Vrn-1g*) were not sufficient to suppress the induction of *COR14b*. In contrast, when plants were grown under long days, the *Vrn-1h* line (low *VRN-1* transcript levels) showed a significantly stronger ( $P= 0.006$ ) induction of *COR14b* than the *Vrn-1f* and *Vrn-1g* mutants (Fig. 4C). As in the previous experiments, both high *VRN-1* transcript levels and long days were necessary for the down-regulation of *COR14b*.

Taken together, the results from this and previous experiments suggest that *VRN-1* expression is required to initiate the developmental processes that reduce the ability of *COR14b* to respond to cold temperatures, but that *VRN-1* transcription alone is not sufficient to produce this effect.

## DISCUSSION

### **Homozygous *mvp/mvp* mutants show more freezing tolerance than *Mvp/-* plants.**

When the shoot apical meristem of the *Mvp/-* plants transitions to the reproductive phase, it stops producing new leaves. Therefore, *Mvp/-* plants are expected to exhibit a reduced ability to generate new leaves after freezing as they transition to the reproductive phase. In contrast the shoot apical meristem of the *mvp/mvp* homozygous mutants never transitions to the reproductive phase and thus, their ability to generate new leaves after freezing should not be drastically altered with time.

While changes in the susceptibility of the shoot apical meristem to freezing damage may account for the observed differences in regrowth after freezing, the *mvp/mvp* homozygous mutants also exhibited greater freezing tolerance in the existing leaves after acclimation than the *Mvp/-* plants as suggested by higher  $F_v/F_m$  values and lower relative conductivity after the freezing treatment. This suggests that the presence of *VRN-1* transcript levels

and the concomitant dampening of *COR14b* induction by cold contributed to a reduction of freezing tolerance in the existing leaves of the *Mvp*<sup>-</sup> plants.

Barley COR14b and the related Arabidopsis COR15 are hydrophilic proteins targeted to the chloroplast stromal compartment (Crosatti et al. 1995). In Arabidopsis, increasing the levels of COR15 results in increased freezing tolerance (Artus et al. 1996). COR15 appears to stabilize membranes from freeze-induced injury, which would account for the reduced electrolyte leakage and higher  $F_v/F_m$  values (Artus et al. 1996). In addition, the higher transcript levels of dehydrin genes such as *DHN5* (Fig. 3) in the leaves of the *mvp/mvp* homozygous mutants relative to the *Mvp*<sup>-</sup> plants likely contributes to their improved freezing tolerance. Dehydrins have a highly conserved 15 amino acid segment (the “K-segment”) which interacts with acidic phospholipids in lipid vesicles. This interaction results in a conformational change of the protein structure that is hypothesized to stabilize membrane integrity (Koag et al. 2003, 2009).

**The increased freezing tolerance of the *mvp* mutants is likely caused by the *VRN-1* deletion.**

The MADS-box meristem identity gene *VRN-1* plays an essential role in the regulation of the transition between vegetative and reproductive phases, and its deletion results in plants that fail to flower. In Arabidopsis, gene duplications of the *VRN-1* homolog that occurred after the monocot-dicot divergence, resulted in three paralogous genes *APETALA1* (*API*), *CAULIFLOWER* (*CAL*), and *FRUITFULL* (*FRU*) that have retained partial ability to promote the transition of the vegetative shoot apical meristem (SAM) to the reproductive phase. Simultaneous deletions of all three genes are required to generate non-flowering Arabidopsis plants (Ferrandiz et al., 2000).

An additional difference between the temperate cereals and Arabidopsis meristem identity genes is their spatial expression profile. *VRN-1* transcripts are detected at high levels in the leaves of wheat (Danyluk et al., 2003; Yan et al., 2003; Li and Dubcovsky, 2008), barley (Schmitz et al., 2000; Trevaskis et al., 2003), *Lolium* (Petersen et al., 2004) and oat (Preston and Kellogg, 2008), suggesting a similar profile among the temperate cereals. In contrast, the Arabidopsis meristem identity homologs are expressed primarily

in the apical meristem and reproductive tissues. *API* and *CAL* transcripts are abundant in the induced SAM and floral primordia in Arabidopsis, but are undetectable or are present at much lower levels in some vegetative tissues such as the vascular tissues of cotyledons (Abe et al., 2005). *FUL* is also expressed primarily in the meristem and floral tissue, but it is also detected in cauline leaves (Teper-Bamnolker and Samach, 2005). In the winter cereals, the expression of *VRN-1* in the leaves (and apices) occurs only after vernalization providing a potentially useful regulatory signal to down-regulate the cold acclimation pathway in this tissue in the spring.

Winter wheat lines exposed to continuous cold temperatures improve their freezing tolerance during the first 3-4 weeks of the treatment but then gradually start losing those gains. The inflexion point in this freezing tolerance curve coincides with the transition of the shoot apical meristem to the double ridge stage and high *VRN-1* transcript levels in the leaves (Danyluk et al., 2003). Limin and Fowler (2006) found that wheat near-isogenic lines for *VRN-1* carrying the allele for winter growth habit tolerate freezing temperatures 11°C lower than lines carrying the *VRN-1* allele for spring growth habit. The authors also showed that when the same near-isogenic spring lines were grown under short days, which are less promotive of *VRN-1* expression than long days, the plants tolerated temperatures 8.5°C colder than the same lines grown under long days (Limin and Fowler, 2006). Based on these results these authors hypothesized that the expression of *VRN-1* might be an important signal to regulate the freezing tolerance pathway.

Previous studies using near-isogenic lines (Limin and Fowler, 2006), and others using QTL mapping for freezing tolerance, all point to the *VRN-1* region as important in the regulation of freezing tolerance (Sutka and Snape, 1989; Roberts, 1990; Hayes et al., 1993; Francia et al., 2004; Galiba et al., 2009). However, the recombination points flanking the *VRN-1* gene in the lines used in these studies are not known and can encompass large chromosome regions including a large number of genes. The use of deletion mutants in this study provides a more precise delimitation of the chromosome region responsible for the differences in freezing tolerance.

Nonetheless, some caution is still required in the interpretation of the *mvp* results, since the deletions present in these lines include other genes flanking *VRN-1*. A screening with

probes for multiple genes in the *VRN-1* region showed that the *mvp* deletions include the *AGLG1*, *CYS*, and *PHYC* genes but exclude flanking genes *CYB5* and *ADA2* (Distelfeld and Dubcovsky, 2010). The exclusion of *ADA2* from the deleted regions is relevant because this gene is critical for *CBF* mediated trans-activation (Stockinger et al., 2001). Based on colinearity with *Brachypodium* the *mvp* deletion is also predicted to include two additional genes, an oligopeptide transporter (*Bradi1g08420*), and a proteinase inhibitor I9 (*Bradi1g08450*) (Distelfeld and Dubcovsky, 2010). Most of these additional genes are unlikely candidates for the improved freezing tolerance observed in the *mvp* mutants, with the exception of phytochrome *PHYC*, since phytochromes B and D have been shown to affect the *CBF* regulon in *Arabidopsis* (Franklin and Whitelam, 2007). Thus, we cannot rule out the possibility of the presence of additional genes with an effect on freezing tolerance in the *mvp* deleted region until it is completely sequenced.

However, the expression studies provided an independent source of evidence pointing to *VRN-1* as the best candidate gene for the down-regulation of the cold acclimation response. In all the *T. monococcum* accessions carrying the *VRN-1f* or *VRN-1g* alleles, the higher transcript levels of *VRN-1* were always associated with a significant down-regulation of *COR14b* when plants were grown in long day conditions (Fig. 4B and 4C, LD). In addition, larger differences in *VRN-1* transcript levels during development were correlated with larger differences in *COR14b* transcript levels (Fig. 4B and 4C, LD). All the expression results presented here support the hypothesis that *VRN-1* is the best candidate for the dampening of the cold acclimation response among the genes present within the *mvp* deletions. We are currently developing TILLING mutants of the *VRN-A1* and *VRN-B1* genes in tetraploid wheat (Uauy et al. 2009) to provide an independent validation of this hypothesis.

### **Homozygous *mvp* mutants exhibit higher transcript levels of several *CBF* and *COR* genes after a short cold treatment**

A negative association between *VRN-1* and *COR* genes *COR14b* and *DHN5* (= *WCS120*) transcript levels has also been reported in previous wheat and barley studies (Vágújfalvi et al., 2000; Danyluk et al., 2003; Knox et al., 2008). In a doubled-haploid barley

population segregating for *VRN-H1*, the lines carrying the recessive *vrn-H1* allele showed higher transcript levels of *CBF* and *COR* genes than those carrying the dominant *Vrn-H1* allele (Stockinger et al., 2007). In addition, lines grown under short days (reduced *VRN-H1* levels) showed higher *CBF* and *COR* transcript levels than lines grown under long days when plants were transferred to the cold (Stockinger et al., 2007). The reductions in the transcript levels of multiple *CBF* genes and their downstream *COR* gene targets in plants with high *VRN-1* transcript levels provide a simple explanation for the gradual decrease in freezing tolerance observed after the initiation of the reproductive phase.

The *mvp* mutants characterized in this study exhibited a similar negative association between *VRN-1* and both *CBF* and *COR* transcription profiles. Five of the eleven *CBF* genes tested by quantitative RT-PCR and the *COR14b* gene showed significantly higher transcript levels 8h after moving the plants to 4°C in the *mvp* deletion homozygotes than in those carrying at least one functional copy of *VRN-1* (Fig. 1 and Fig. S4). In plants maintained at 4°C for 12 days, the *COR14b* transcript levels were still 10-fold higher in the plants homozygous for the *mvp* deletion (Fig. 2). However, analyses of two additional *COR* genes, *DHN5* and *DHN8* showed that not all *COR* genes respond in the same way. Whereas *DHN5* showed the same negative correlation with *VRN-1* as *COR14b*, the *DHN8* gene was not significantly affected by the change in the level of *VRN-1* transcripts, which indicates that not all *COR* genes are down-regulated by *VRN-1*. Similarly, for half of the 11 *CBF* genes present in the *FR-2* cluster, no significant differences in transcript levels were detected between *Mvp*<sup>-/-</sup> and *mvp/mvp* lines (Fig. 1 and Fig. S4), suggesting that cold activation of these genes is not regulated by *VRN-1*. Thus it appears that this *VRN-1*-mediated mechanism may play a role in the regulation of a specific subset of cold responsive genes.

**Allelic differences in *VRN-1* are likely sufficient to explain differences in freezing tolerances previously assigned to a separate *FR-1* locus.**

In earlier studies, differences in freezing tolerance mapped to the *VRN-1* region in wheat were considered to be the result of a closely linked gene designated *FR-1*. However, only two studies have reported recombination between *FR-1* and *VRN-1*, and they differ in the

relative position of these two genes, with *FR-1* distal to *VRN-1* in the initial mapping studies (Galiba et al. 1995, 1997) and proximal to *VRN-1* in a later mapping study using deletion lines (Sutka et al., 1999). Although the differences in freezing tolerance across the deletion lines used in the latter study were clear, it is still possible that the reduced freezing tolerance observed in the larger deletion used to map *FR-1* to a proximal deletion bin than *VRN-1*, was the result of the loss of a larger number of genes and an overall reduction in plant vigor rather than the effect of a single *FR-1* gene. It is also possible that simultaneous segregation at the linked *FR-2* locus, which was not known at the time of these two studies, affected the mapping results.

The improved freezing tolerance and higher transcript levels of *CBF* and *COR* genes in the *mvp* mutants suggests that *VRN-1* allelic differences are likely sufficient to explain differences in freezing tolerance previously considered to be the result of a separate *FR-1* gene. Therefore, our results support the hypothesis that *FR-1* is a pleiotropic effect of *VRN-1* rather than a separate gene. This hypothesis is also supported by experiments showing that the repression of *VRN-1* by short days in spring wheat genotypes is associated with increased freezing tolerance (Limin et al., 2006), and that *VRN-1* transcript levels in the different Triple Dirk near-isogenic lines are inversely correlated with freezing tolerance (Koemel et al., 2004).

### ***VRN-1* transcription is not sufficient to promote the down-regulation of *COR* genes**

The experiment using the *T. monococcum* lines carrying the *VRN-1f* and *VRN-1g* alleles (Fig. 4) shows that up-regulation of *VRN-1* transcript levels under short days is insufficient to produce a significant down-regulation of *COR14b* as that observed under long days. Under short days, six-week old plants carrying these alleles show high transcript levels of *VRN-1* and a transition of the vegetative apex to the double ridge stage. However, under continuous short days spike development proceeds slowly and stems fail to elongate. Once plants are transferred to long days, genotypes having the *VRN-1f* and *VRN-1g* alleles complete their spike development faster and head earlier than genotypes with the wild-type *VRN-1* allele due to their more advanced developmental state (Dubcovsky et al., 2006).

When grown under long days, plants carrying the *VRN-1f* and *VRN-1g* alleles showed significantly lower levels of *COR14b* than lines carrying the *vrn-1* or *Vrn-1h* alleles. These results were consistent across three independent accessions for each of the *VRN-1* alleles supporting the hypothesis that the differences in *COR14b* were associated to the differences in the *VRN-1* alleles. However, no differences in *COR14b* were observed among the same genotypes under short days despite large differences in *VRN-1* transcript levels. These results suggest that the down-regulation of the *COR14b* requires the presence of additional factors that are activated under long days and that require the expression of *VRN-1*. Taken together, the *mvp* mutant and *VRN-1f* and *VRN-1g* experiments suggest that *VRN-1* expression is necessary but not sufficient to down-regulate several *COR* genes and reduce freezing tolerance in the leaves of wheat plants. A similar phenomenon has recently been described in Arabidopsis where the floral activator MADS-box gene *SOC1* functions as a negative regulator of the cold response pathway through the direct repression of the *CBF* genes (Seo et al., 2009). In Columbia wild type, *SOC1* was expressed most strongly in leaves, but was also detected in vegetative apices, inflorescence, stems of flowering plants, and roots (Lee et al., 2000). A microarray experiment comparing seven-day old seedlings of a *soc1* knock-out mutant and a *SOC-1* over-expressing line with wild type plants revealed that six *COR* genes were amongst the 20 genes most negatively regulated in the *SOC-1* over-expressing line. In addition, the expression level of the three Arabidopsis *CBF* genes increased in the *soc1* mutants and decreased in *SOC-1* over-expression lines, without affecting the transcript level of the *CBF* regulatory genes *ICE1*, *HOS1* or *ZAT12*. A chromatin immunoprecipitation (ChIP) experiment using a *SOC1* antibody revealed that the CArG-box regions in the *CBF* promoters were enriched in the *SOC1* over-expressing line relative to the *soc1* knock-out, which suggests that *SOC1* negatively regulates cold response through direct repression of the transcription of the *CBF* genes (Seo et al., 2009). It is interesting to point out here that the expression of the wheat homolog of Arabidopsis *SOC1*, *WSOC1*, is not affected by vernalization or photoperiod, suggesting different functions in these two species (Shitsukawa et al., 2007b).

Although both *SOC1* in Arabidopsis and *VRN-1* in the temperate cereals seem to be associated with the down-regulation of the *CBF* and *COR* genes in the leaves, the effect

of *VRN-1* on the *CBF* genes does not seem to be as direct as the effect of *SOC1* in Arabidopsis. The results from the experiments using *T. monococcum* accessions with differential expression of *VRN-1* under short days suggest that additional genes operating downstream of *VRN-1* and that are regulated by long-days are required to mediate the negative effect of *VRN-1* on freezing tolerance. The identification of these downstream genes and the understanding of their regulatory mechanisms could potentially lead to novel strategies to prevent the premature dampening of the cold acclimation pathway in environments where the premature activation of *VRN-1* may increase the risk of freezing damage.

### **Do temperate cereals respond differently to the same cool temperatures in the fall and the spring?**

The system described above provides the temperate cereals with the ability to differentiate the same cool temperature in the fall and the spring. A cool temperature in the fall, when plants have low *VRN-1* transcript levels, results in the induction of the *CBF* and the downstream *COR* genes initiating the acclimation of the plants to cold temperatures. This is essential in the fall, when cool temperatures are an indication of the approaching freezing temperatures of the winter. The same cool temperature in the spring, when *VRN-1* transcript levels in the leaves increase significantly in response to lengthening photoperiod, would result in a significantly lower up-regulation of several *CBF* and *COR* genes. Since cool temperatures in spring are generally not a prelude of coming freezing temperatures, a robust up-regulation of the CBF pathway response in the spring would likely not be advantageous for plant survival.

A similar system seems to be operating in Arabidopsis. Arabidopsis *soc1* null mutants show increased responsiveness of the *CBF* genes to cold and improved freezing tolerance suggesting that low levels of *SOC1* transcripts during the fall may favor plant acclimation to cold temperatures (Seo et al., 2009). *SOC1* transcript levels increase significantly by the time of the initiation of Arabidopsis flowering (Lee et al., 2000), indicating that high *SOC1* transcript levels will be present in the leaves in the spring, down-regulating the *CBF* genes and their downstream *COR* targets.

The activation of the *CBF*-regulon has a potentially high energetic cost to plants since numerous *COR* genes are up-regulated in the leaves by these transcription factors (Fowler and Thomashow, 2002). In addition, *CBF* genes have also been shown to repress plant growth (Achard et al. 2008). Therefore, the down-regulation of the *CBF* genes during the spring has a potential adaptative value, ensuring the plant's rapid development under optimal conditions and may explain the presence of related systems in Arabidopsis and the temperate cereals.

## MATERIALS AND METHODS

### Plant materials

**Maintained vegetative phase (*mvp*) mutants:** Two independent *T. monococcum* mutants (*mvp-1* and *mvp-2*) that remain indefinitely in the vegetative state (Shitsukawa et al., 2007a) were used in this study. Since the two mutants carry similar deletions (Distelfeld and Dubcovsky, 2010), they were alternated among experiments depending on seed supply. Seeds from these mutants were kindly provided by K. Murai. These mutants were generated by ion-beam radiation and both have large deletions that include *VRN-1* (Shitsukawa et al., 2007a) and several flanking genes (Distelfeld and Dubcovsky, 2010). The *mvp-1* mutation was generated in the KU104-2 background, and the *mvp-2* mutation in the KU104-1 background (Shitsukawa et al., 2007a). When grown in the greenhouse under long day conditions (16 h photoperiods) KU104-2 flowered 10 weeks after planting while KU104-1 flowered three weeks later.

Homozygous *mvp* individuals do not flower and therefore must be maintained in a heterozygous state. Genotyping was carried out using a dominant *VRN-1* molecular marker based on a set of three primers that are described in the supporting online materials (Fig. S1). Using this assay the lines carrying one or two functional *VRN-1* copies are detected as a single genotypic class, referred throughout the text as “*Mvp*/-”.

***VRN-1f*, *VRN-1g* and *VRN-1h* alleles:** *Triticum monococcum* lines with four different *VRN-1* alleles were used to test the effect of their differential regulation under short days on *COR14b* transcript levels. *Triticum monococcum* lines PI355546, PI427927, and

PI237659 carry the “wild type” *vrn-1* allele and a recessive *vrn-2* allele that confers spring growth habit. Lines with the *vrn-1* allele showed no expression under short days in previous studies (Dubcovsky et al., 2006). *T. monococcum* lines PI191097, PI192063, and PI393496 carry the *Vrn-1f* allele, which has a 1-bp deletion in a CArG box located in the promoter, plus an insertion of a repetitive element in the first intron. *T. monococcum* lines PI326317, PI418582, and PI349049 carry the *Vrn-1g* allele, which has a 34-bp deletion including the promoter CArG box. Both the *Vrn-1f* and *Vrn-1g* alleles confer spring growth habit and show high levels of *VRN-1* transcripts under short days (Dubcovsky et al., 2006). *T. monococcum* accession PI306540 has the *VRN-1h* allele that has the same intron one insertion as *Vrn-1f* but lacks the CArG box mutation. This allele is not expressed under short days and confers a spring phenotype but with later flowering than the *Vrn-1f* and *Vrn-1g* alleles (Dubcovsky et al., 2006). Although the shoot apical meristem transitions to the double ridge stage, spike development progresses slowly and spikes fail to elongate if these plants are left under short days (Dubcovsky et al., 2006).

## Methods

**Growth, cold acclimation, and freezing assays:** Freezing experiments with the *mvp* mutants were all carried out using long day conditions (LD, 16 h light / 8 h dark). Experiments with the *T. monococcum* lines with different *VRN-1* alleles were carried out using both long days and short day (SD, 8 h light / 16 h dark) photoperiod cycles.

The RNA blot analysis and the freezing experiment of the *mvp-2* mutants were done in parallel. Seed collected from *Mvp-2*<sup>-/-</sup> heterozygotes were grown under cool white fluorescent lamps in the laboratory at room temperature for 13 days using a light intensity of 50  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Genotyped seedlings were transplanted to wooden boxes having internal dimensions measuring 42 cm (l) x 22 cm (w) x 14 cm (h) and having 9.5 cm soil depth. The boxes were placed into a Conviron growth chamber (model PGW36, Controlled Environments Inc., ND) for an additional 47 d under cool white fluorescent and incandescent bulbs, using a light intensity of 130  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , and a constant temperature of 23 °C. The growth chamber temperature was decreased to 10 °C and held at this temperature for 19 days.

Wooden boxes were then transferred to Percival growth chambers (model CU-36L2X, Geneva Scientific, Fontana, WI) where the seedlings were held at 4 °C for 12 d (the PGW36 growth chamber can only cool to +10°C). After cold acclimation the CU-36L2X growth chamber temperatures were decreased to -2 °C, and held for 12 h. Ice nucleation was induced by spraying the leaves with ice water. Following 12 h at -2 °C, the temperature was then decreased 1 °C per h to the target temperatures of -9°C and -11 °C and held at these temperatures for 24 h. Afterwards the temperature was returned to +2 °C for 12 h. During this time the plants were kept in the dark. The chamber temperature was then raised to 20 °C and returned to a 16 h photoperiod.

**Chlorophyll fluorescence ( $F_v/F_m$ ):** Chlorophyll fluorescence measurements were made using a hand-held portable pulse amplitude modulated fluorometer (model OS-30p, Opti Sciences, Hudson, NH).  $F_v/F_m$  measurements were taken 2-4 h after the plants were returned to normal growth conditions. Leaves were dark adapted for approximately 10 minutes prior to taking the measurements.

**Relative conductivity:** Conductivity measurements were taken on crown tissue consisting of a 1 to 1.5 cm segment of the white, nonphotosynthetic tissue between the upper photosynthetic green shoot and the primary root. Because the use of the crowns required the destruction of the plant the number of replications in the first experiment was limited to three *mvp/mvp* homozygotes and six *Mvp/-* plants. The second experiment used leaves instead of crown regions and nine plants from temperature / genotype combination. The electrolyte leakage assay methods are described in detail in the SOM in the section B of ‘Methods used for Figure S3’). Tubes were shaken for 1 h at 300 rpm before reading the conductivity with an Accumet Basic AB30 electrical conductivity meter (Fisher Scientific). Tubes were then autoclaved for 20 minutes, cooled to room temperature and shaken for 1 h at 300 rpm before measuring the total potential conductivity. Values were adjusted by subtracting the conductivity of the deionized water. Relative conductivity represents the adjusted mean ion leakage as a percentage of the total adjusted leakage from frozen-killed samples (see formula in SOM).

**CBF qRT-PCR experiments:** Eight *mvp-2/mvp-2* and *Mvp-2/-* and ten *mvp-1/mvp-1* and *Mvp-1/-* plants were selected using the *VRN-1* molecular marker and were grown in the

greenhouse for four weeks and eight weeks, respectively (20-25 °C, long days). Plants were then transferred to a growth chamber at 4 °C for 8 hours. RNA samples were collected from leaves from 8 *mvp-2* and 8 *Mvp-2*<sup>-/-</sup> plants before (20 °C) and after (4 °C) the cold treatment in the first experiment, and from 10 *mvp-1* and 10 *Mvp-1*<sup>-/-</sup> plants only after the cold treatment in the second experiment.

***COR14b* qRT-PCR time course experiment:** Eight *mvp-2* and eight *Mvp-2*<sup>-/-</sup> plants were selected using the *VRN-1* molecular markers and grown in the greenhouse under the same conditions described above. After four weeks, when the *Mvp-2*<sup>-/-</sup> plants were still at the vegetative stage, plants were transferred to 4 °C and were kept at this temperature for 12 days at the same light intensity indicated above (long days). Leaf samples for RNA analysis were collected one day before the cold treatment and after 8 h, 32 h, 4 days and 12 days at 4 °C. Samples were always collected at 2:00 pm (8 h after the subjective daybreak) to avoid potential differences at different times of the day.

***VRN-1/COR14b* qRT-PCR experiment:** Five plants with each *VRN-1* allele were grown for either two or six weeks under short or long day conditions before transferring to 4 °C. RNA samples were collected from leaves after 32 hours of cold treatment.

For all qRT-PCR experiments, RNA was extracted using the Spectrum™ Plant Total RNA Kit (Sigma-Aldrich, Saint Louis, MO). First strand cDNA was synthesized from 1 µg of total RNA using the QuantiTect Reverse Transcription Kit (Qiagen, Valencia, CA). Primers for qRT-PCR expression analyses are presented in Supplementary Table S1. The *TRANSLATION ELONGATION FACTOR 1 (TEF1)* and *ACTIN* genes were used as endogenous controls (Distlefeld and Dubcovsky 2010).

***RNA blot analyses:*** Tissue samples were collected 13-14 h after the subjective daybreak from both cold-acclimating plants and non-acclimated plants. Total RNA was isolated using RNeasy Plant Mini kits (Qiagen, Valencia, CA). Seven µg of total RNA was loaded per lane. RNA samples consisted of RNAs pooled from the crown tissue of ten (*mvp-2*<sup>-/-</sup> experiment) or five (*mvp-1* and *VRN-1* promoter deletion mutant experiments) plants.

Fragments used as probes were generated by PCR amplification of cloned cDNA inserts. Radiolabeled probes were generated by random priming (Feinberg and Vogelstein, 1983). Overnight hybridizations were at 42°C in 50% formamide, 5X SSC, 20 mM Na-

phosphate buffer pH 6.8 containing 100 µg/ml herring sperm DNA, 1X Denhardt's solution, 0.1% SDS, and 10% dextran sulfate. Three to four 1 h moderate-stringency washes were performed at 62-65°C in 0.2X SSC, 0.05% SDS and 0.01% Na-pyrophosphate. Images were generated using the Molecular Dynamics Storm840 PhosphorImager (GE Healthcare) and phosphor autoradiography.

Upon request, all novel materials described in this publication will be made available in a timely manner for non-commercial research purposes, subject to the requisite permission from any third-party owners of all or parts of the material. Obtaining any permission will be the responsibility of the requestor.

## **ACKNOWLEDGEMENTS**

The authors thank Dr. Koji Murai for providing the seeds for the *mvp-1* and *mvp-2* mutants.

## REFERENCES

- Abe M, Kobayashi Y, Yamamoto S, Daimon Y, Yamaguchi A, Ikeda Y, Ichinoki H, Notaguchi M, Goto K, Araki T** (2005) FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science* **309**: 1052-1056
- Achard P, Gong F, Cheminant S, Alioua M, Hedden P, Genschik P** (2008) The cold-inducible CBF1 factor-dependent signaling pathway modulates the accumulation of the growth-repressing DELLA proteins via its effect on gibberellin metabolism. *The Plant Cell* **20**: 2117-2129
- Artus NN, Uemura M, Steponkus PL, Gilmour SJ, Lin C, Thomashow MF** (1996) Constitutive expression of the cold-regulated *Arabidopsis thaliana* *COR15a* gene affects both chloroplast and protoplast freezing tolerance. *Proc Natl Acad Sci USA* **93**: 13404-13409
- Båga M, Chodaparambil SV, Limin AE, Pecar M, Fowler DB, Chibbar RN** (2007) Identification of quantitative trait loci and associated candidate genes for low-temperature tolerance in cold-hardy winter wheat. *Funct Integ Genomics* **7**: 53-68
- Brule-Babel AL, Fowler DB** (1988) Genetic control of cold hardiness and vernalization requirement in winter wheat. *Crop Sci* **28**: 879-884
- Butler WL, Kitajima M** (1975) Fluorescence quenching in photosystem II of chloroplasts. *Biochim Biophys Acta* **376**: 116-125
- Choi DW, Zhu B, Close TJ** (1999) The barley (*Hordeum vulgare* L.) dehydrin multigene family: sequences, allele types, chromosome assignments, and expression characteristics of 11 *Dhn* genes of cv Dicktoo. *Theor Appl Genet* **98**: 1234-1247
- Crosatti C, Nevo E, Stanca AM, Cattivelli L** (1996) Genetic analysis of the accumulation of COR14 proteins in wild (*Hordeum spontaneum*) and cultivated (*Hordeum vulgare*) barley. *Theor Appl Genet* **93**: 975-981
- Crosatti C, Soncini C, Stanca AM, Cattivelli L** (1995) The accumulation of a cold-regulated chloroplastic protein is light-dependent. *Planta* **196**: 458-463

- Dal Bosco C, Busconi M, Govoni C, Baldi P, Stanca AM, Crosatti C, Bassi R, Cattivelli L** (2003) *Cor* gene expression in barley mutants affected in chloroplast development and photosynthetic electron transport. *Plant Physiol.* **131**: 793-802
- Dexter ST** (1956) The evaluation of crop plants for winter hardiness. *Adv Agron* **8**: 203-239
- Danyluk J, Houde M, Rassart E, Sarhan F** (1994) Differential expression of a gene encoding an acidic dehydrin in chilling sensitive and freezing tolerant *gramineae* species. *FEBS Lett* **344**: 20-24
- Danyluk J, Kane NA, Breton G, Limin AE, Fowler DB, Sarhan F** (2003) *TaVRT-1*, a putative transcription factor associated with vegetative to reproductive transition in cereals. *Plant Physiol* **132**: 1849-1860
- Danyluk J, Perron A, Houde M, Limin A, Fowler B, Benhamou N, Sarhan F** (1998) Accumulation of an acidic dehydrin in the vicinity of the plasma membrane during cold acclimation of wheat. *Plant Cell* **10**: 623-638
- Distelfeld A, Dubcovsky J** (2010) Characterization of the *maintained vegetative phase (mvp)* deletions from einkorn wheat and their effect on *VRN2* and *FT* transcript levels. *Mol Genet Genomics* **283**: 223-232.
- Distelfeld A, Li C, Dubcovsky J** (2009) Regulation of flowering in temperate cereals. *Curr Opin Plant Biol* **12**: 178-184
- Dubcovsky J, Loukoianov A, Fu D, Valarik M, Sanchez A, Yan L** (2006) Effect of photoperiod on the regulation of wheat vernalization genes *VRN1* and *VRN2*. *Plant Mol Biol* **60**: 469-480
- Feinberg AP, Vogelstein P** (1983) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* **132**: 6-13
- Ferrandiz C, Gu Q, Martienssen R, Yanofsky MF** (2000) Redundant regulation of meristem identity and plant architecture by *FRUITFULL*, *APETALA1* and *CAULIFLOWER*. *Development* **127**: 725-734

- Fowler DB, Chauvin LP, Limin AE, Sarhan F** (1996a) The regulatory role of vernalization in the expression of low-temperature-induced genes in wheat and rye. *Theor Appl Genet* **93**: 554-559
- Fowler DB, Breton G, Limin AE, Mahfoozi S, Sarhan F** (2001) Photoperiod and temperature interactions regulate low-temperature-induced gene expression in barley. *Plant Physiol* **127**: 1676-1681
- Fowler DB, Limin AE** (2004) Interactions among factors regulating phenological development and acclimation rate determine low-temperature tolerance in wheat. *Ann Bot-London* **94**: 717-724
- Fowler DB, Limin AE, Ritchie JT** (1999) Low-temperature tolerance in cereals: Model and genetic interpretation. *Crop Sci* **39**: 626-633
- Fowler DB, Limin AE, Wang SY, Ward RW** (1996b) Relationship between low-temperature tolerance and vernalization response in wheat and rye. *Can J Plant Sci* **76**: 37-42
- Fowler S, Thomashow MF** (2002) Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell* **14**: 1675-1690
- Francia E, Barabaschi D, Tondelli A, Laidò G, Rizza F, Stanca AM, Busconi M, Fogher C, Stockinger EJ, Pecchioni N** (2007) Fine mapping of a *HvCBF* gene cluster at the frost resistance locus *Fr-H2* in barley. *Theor Appl Genet* **115**: 1083-1091
- Francia E, Rizza F, Cattivelli L, Stanca AM, Galiba G, Tóth B, Hayes PM, Skinner JS, Pecchioni N** (2004) Two loci on chromosome 5H determine low-temperature tolerance in a Nure (winter) × Tremois (spring) barley map. *Theor Appl Genet* **108**: 670-680
- Franklin KA, Whitelam GC** (2007) Light-quality regulation of freezing tolerance in *Arabidopsis thaliana*. *Nature Genetics* **39**: 1410-1413
- Fu D, Szúcs P, Yan L, Helguera M, Skinner J, Hayes P, Dubcovsky J** (2005) Large deletions in the first intron of the *VRN-1* vernalization gene are associated with

- spring growth habit in barley and polyploid wheat. *Mol Genet Genomics* **273**: 54-65
- Galiba G, Quarrie SA, Sutka J, Morgounov A** (1995) RFLP mapping of the vernalization (*Vrn1*) and frost resistance (*Fr1*) genes on chromosome 5A of wheat. *Theor Appl Genet* **90**: 1174-1179
- Galiba G, Kerepsi I, Snape JW, Sutka J** (1997) Location of a gene regulating cold-induced carbohydrate production on chromosome 5A of wheat. *Theor Appl Genet* **95**: 265-270
- Galiba, G, Vágújfalvi, A, Li, C, Soltész, A, Dubcovsky, J** (2009) Regulatory genes involved in the determination of frost tolerance in temperate cereals. *Plant Sci* **176**: 12–19
- Grossi M, Giorni E, Rizza F, Stanca AM, Cattivelli L** (1998) Wild and cultivated barleys show differences in the expression pattern of a cold-regulated gene family under different light and temperature conditions. *Plant Mol Biol* **38**: 1061-1069
- Hayes HK, Aamodt OS** (1927) Inheritance of winter hardiness and growth habit in crosses of Marquis with Minhardi and Minturki wheats. *J Agric Res* **35**: 223-236
- Hayes PM, Blake T, Chen THH, Tragoonrung S, Chen F, Pan A, Liu B** (1993) Quantitative trait loci on barley (*Hordeum vulgare* L.) chromosome 7 associated with components of winterhardiness. *Genome* **36**: 66-71
- Houde M, Dhindsa RS, Sarhan F** (1992) A molecular marker to select for freezing tolerance in Gramineae. *Mol Gen Genet* **234**: 43-48.
- Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF** (1998) Arabidopsis *CBF1* overexpression induces *COR* genes and enhances freezing tolerance. *Science* **280**: 104-106
- Koag MC, Fenton RD, Wilkens S, Close TJ** (2003) The binding of maize DHN1 to lipid vesicles. Gain of structure and lipid specificity. *Plant Physiol* **131**: 309-316

- Koag MC, Wilkens S, Fenton RD, Resnik J, Vo E, Close TJ** (2009) The K-segment of maize DHN1 mediates binding to anionic phospholipid vesicles and concomitant structural changes. *Plant Physiol* **150**: 1503-1514
- Koemel JE, Jr., Guenzi AC, Anderson JA, Smith EL** (2004) Cold hardiness of wheat near-isogenic lines differing in vernalization alleles. *Theor Appl Genet* **109**: 839-846
- Knox AK, Li C, Vágújfalvi A, Galiba G, Stockinger EJ, Dubcovsky J** (2008) Identification of candidate *CBF* genes for the frost tolerance locus *Fr-A<sup>m</sup>2* in *Triticum monococcum*. *Plant Mol Biol* **67**: 257-270
- Knox AK, Dhillon T, Cheng H, Tondelli A, Pecchioni N, Stockinger EJ** (2010) *CBF* gene copy number variation at *Frost Resistance-2* is associated with levels of freezing tolerance in temperate-climate cereals. *Theor Appl Genet* **121**: 21-35.
- Lee H, Suh SS, Park E, Cho E, Ahn JH, Kim SG, Lee JS, Kwon YM, Lee I** (2000) The AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in Arabidopsis. *Genes Dev* **14**: 2366-2376
- Li C, Dubcovsky J** (2008) Wheat FT protein regulates *VRN1* transcription through interactions with FDL2. *Plant J* **55**: 543-554
- Limin AE, Danyluk J, Chauvin LP, Fowler DB, Sarhan F** (1997) Chromosome mapping of low-temperature induced *Wcs120* family genes and regulation of cold-tolerance expression in wheat. *Mol Gen Genet* **253**: 720-727
- Limin AE, Fowler DB** (2006) Low-temperature tolerance and genetic potential in wheat (*Triticum aestivum* L.): response to photoperiod, vernalization, and plant development. *Planta* **224**: 360-366
- Livak KJ, Schmittgen TD** (2001) Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta C_T}$  method. *Methods* **25**: 402-408
- Loukoianov A, Yan L, Blechl A, Sanchez A, Dubcovsky J** (2005) Regulation of *VRN-1* vernalization genes in normal and transgenic polyploid wheat. *Plant Physiol* **138**: 2364-2373

- Miller AK, Galiba G, Dubcovsky J** (2006) A cluster of 11 *CBF* transcription factors is located at the frost tolerance locus *Fr-A<sup>m</sup>2* in *Triticum monococcum*. *Mol Genet Genomics* **275**: 193-203
- NDong C, Danyluk J, Wilson KE, Poccock T, Huner NP, Sarhan F** (2002) Cold-regulated cereal chloroplast late embryogenesis abundant-like proteins. Molecular characterization and functional analyses. *Plant Physiol* **129**: 1368-1381.
- Petersen K, Didion T, Andersen CH, Nielsen KK** (2004) MADS-box genes from perennial ryegrass differentially expressed during transition from vegetative to reproductive growth. *J Plant Physiol* **161**: 439-447
- Pidal B, Yan L, Fu D, Zhang F, Tranquilli G, Dubcovsky J** (2009) The *CArG*-box in the promoter region of wheat vernalization gene *VRNI* is not necessary to mediate the vernalization response. *J Hered* **100**: 355-364
- Preston JC, Kellogg EA** (2008) Discrete developmental roles for temperate cereal grass *VRNI/FUL*-like genes in flowering competency and the transition to flowering. *Plant Physiol* **146**: 265-276
- Roberts DWA** (1986) Chromosomes in Cadet and Rescue wheats carrying loci for cold hardiness and vernalization response. *Can J Genet Cytol* **28**: 991-997
- Roberts DWA** (1990) Identification of loci on chromosome-5A of wheat involved in control of cold hardiness, vernalization, leaf length, rosette growth habit, and height of hardened plants. *Genome* **33**: 247-259
- Schmitz J, Franzen R, Ngyuen TH, Garcia-Maroto F, Pozzi C, Salamini F, Rohde W** (2000) Cloning, mapping and expression analysis of barley MADS-box genes. *Plant Mol Biol* **42**: 899-913
- Seo E, Lee H, Jeon J, Park H, Kim J, Noh YS, Lee I** (2009) Crosstalk between cold response and flowering in *Arabidopsis* is mediated through the flowering-time gene *SOCl* and Its upstream negative regulator *FLC*. *Plant Cell* **21**: 3185-3197
- Shitsukawa N, Ikari C, Shimada S, Kitagawa S, Sakamoto K, Saito H, Ryuto H, Fukunishi N, Abe T, Takumi S, Nasuda S, Murai K** (2007a) The einkorn

- wheat (*Triticum monococcum*) mutant, *maintained vegetative phase*, is caused by a deletion in the *VRN1* gene. *Genes Genet Syst* **82**: 167-170
- Shitsukawa N, Ikari C, Mitsuya T, Sakiyama T, Ishikawa A, Takumi S, Murai K** (2007b) Wheat *SOC1* functions independently of *WAP1/VRN1*, an integrator of vernalization and photoperiod flowering promotion pathways. *Physiologia Plantarum* **130**: 627-636
- Skinner J, Szűcs P, von Zitzewitz J, Marquez-Cedillo L, Filichkin T, Stockinger EJ, Thomashow MF, Chen THH, Hayes PM** (2006) Mapping of barley homologs to genes that regulate low temperature tolerance in Arabidopsis. *Theor Appl Genet* **112**: 832-842
- Stockinger EJ, Gilmour SJ, Thomashow MF** (1997) Arabidopsis thaliana *CBF1* encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc Natl Acad Sci USA* **94**: 1035-1040
- Stockinger EJ, Mao YP, Regier MK, Triezenberg SJ, Thomashow MF** (2001) Transcriptional adaptor and histone acetyltransferase proteins in Arabidopsis and their interactions with CBF1, a transcriptional activator involved in cold-regulated gene expression. *Nucleic Acids Res* **29**: 1524-1533
- Stockinger EJ, Skinner JS, Gardner KG, Francia E, Pecchioni N** (2007) Expression levels of barley *Cbf* genes at the *Frost resistance-H2* locus are dependent upon alleles at *Fr-H1* and *Fr-H2*. *Plant J* **51**: 308-321
- Sutka J, Snape JW** (1989) Location of a gene for frost resistance on chromosome 5A of wheat. *Euphytica* **42**: 41-44
- Sutka J, Galiba G, Vagujfalvi A, Gill BS, Snape JW** (1999) Physical mapping of the *Vrn-A1* and *Fr1* genes on chromosome 5A of wheat using deletion lines. *Theor Appl Genet* **99**: 199-202

- Teper-Bamnolker P, Samach A** (2005) The flowering integrator FT regulates SEPALLATA3 and FRUITFULL accumulation in Arabidopsis leaves. *Plant Cell* **17**: 2661-2675
- Thomashow MF** (1990) Molecular genetics of cold acclimation in higher plants. *Adv Genet* **28**: 99-131
- Thomashow MF** (1999) PLANT COLD ACCLIMATION: Freezing tolerance genes and regulatory mechanisms. *Annu Rev Plant Phys* **50**: 571-599
- Trevaskis B, Bagnall DJ, Ellis MH, Peacock WJ, Dennis ES** (2003) MADS box genes control vernalization-induced flowering in cereals. *Proc Natl Acad Sci USA* **100**: 13099-13104
- Trevaskis B, Hemmin MN, Dennis ES, Peacock WJ** (2007) The molecular basis of vernalization-induced flowering in cereals. *Trends Plant Sci* **12**: 352-357
- Uauy C, Paraiso F, Colasuonno P, Tran RK, Tsai H, Berardi S, Comai L, Dubcovsky J** (2009) A modified TILLING approach to detect induced mutations in tetraploid and hexaploid wheat *BMC Plant Biol* **9**: 115-128
- Vágújfalvi A, Crosatti C, Galiba G, Dubcovsky J, Cattivelli L** (2000) Two loci on wheat chromosome 5A regulate the differential cold-dependent expression of the *cor14b* gene in frost-tolerant and frost-sensitive genotypes. *Mol Gen Genet* **263**: 194-200
- Vágújfalvi A, Galiba G, Cattivelli L, Dubcovsky J** (2003) The cold regulated transcriptional activator *Cbf3* is linked to the frost-tolerance gene *Fr-A2* on wheat chromosome 5A. *Mol Genet Genomics* **269**: 60-67
- Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF** (2005) Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of Arabidopsis. *Plant J.* **41**: 195-211
- Yan L, Helguera M, Kato K, Fukuyama S, Sherman J, Dubcovsky J** (2004) Allelic variation at the *VRN-1* promoter region in polyploid wheat. *Theor Appl Genet* **109**: 1677-1686

**Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J (2003)**  
Positional cloning of wheat vernalization gene *VRN1*. Proc Natl Acad Sci USA  
**100:** 6263-6268

## Figure legends

**Fig.1:** Quantitative RT-PCR analysis of transcript levels of the *CBF* genes present at the *FR-2* locus relative to the *ACTIN* endogenous control. Samples were collected from leaves of 4-week old *mvp-2/mvp-2* and *Mvp-2/-* plants (20 °C) and again one day later at the same time following an 8 hour cold treatment at 4 °C. Values in the Y axes were normalized and calibrated using the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen 2001). The same calibrator was used for all genes so scales are comparable across genes. Values are averages of 8 biological replications  $\pm$  SE of the means. The inset shows *CBF14* transcript levels which were significantly higher than the other genes at this locus. *P* values for the differences between *mvp/mvp* and *Mvp/-* after the cold treatment were calculated using ANOVA and are indicated by \*:  $P < 0.05$  or \*\*:  $P < 0.01$ ).

**Fig. 2.** Quantitative RT-PCR transcript levels of *COR14b* relative to *TEF1* endogenous control. Plants were 4 weeks old at the beginning of the experiment and were exposed to 4 °C for 12 days. Values in the Y axes are normalized and calibrated values using the  $2^{\Delta\Delta CT}$  method (Livak and Schmittgen 2001). Homozygous *mvp-2/ mvp-2* plants (null *VRN-1*) are indicated by black squares and lines and *Mvp-2/-* plants (1 or 2 *VRN-1* copies) by gray triangles and lines. Values are averages of 8 biological replications in the untransformed scale  $\pm$  SE of the means. *P* values were calculated using ANOVA of  $\log(n+1)$  transformed values for each time point (\*\*:  $P < 0.01$ ).

**Fig. 3.** Time courses of *VRN-1*, and *COR* genes *COR14b*, *DHN5*, and *DHN8* transcripts in *mvp-2/ mvp-2* plants homozygous for a *VRN-1* deletion (*m*) and *Mvp-2/-* plants with one or two functional *VRN-1* copies (*M*). The presence of a faint *VRN-1* hybridization in the 87 days mutant sample is suspected to be due to cross contamination.


**Fig. 4.** *VRN-1* and *COR14b* transcript levels in a set of *T. monococcum* lines differing in *VRN-1* expression under short days. Abbreviations: W= warm conditions, C= decrease from 18 °C to 6 °C (occurring at daybreak), 2W= two-week old, 6W= six-week old, SD = short day, LD = long day, wt= wild type, *Vrn-A1f*= allele with 1-bp deletion in the CArG box coupled with *VRN-1* intron 1 insertion, *Vrn-A1g* = allele with 34-bp deletion encompassing the CArG box. *Vrn-A1h* = allele with insertion in *VRN-1* intron 1. The three accessions with the *Vrn-A1f* allele carry the dominant *Vrn-A2* allele whereas all the other accessions carry non-functional *vrn-A2* alleles. All the accessions have a spring growth habit. A) mRNA blot analyses of three genotypes per promoter class (indicated by different PI numbers). Arrows and arrowheads identify the presence and absence, respectively, of *COR14b* transcripts in 6W plants in the *Vrn-A1f* and *Vrn-A1g* natural mutants under SD and LD. B-C) qRT-PCR validation of *VRN-1* and *COR14b* transcript levels relative to *ACTIN* endogenous control at 4 °C. Values in the Y axes are normalized and calibrated values using the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen 2001). Lines carrying the wild type allele (*vrn-1*) or the *Vrn-1h* (spring, not induced at SD) are indicated by black bars and lines carrying the *Vrn-1f* and *Vrn-1g* alleles (spring, induced at SD) by gray bars. Values are averages of 5 biological replications  $\pm$  SE of the means. *P* values were calculated using contrasts between either *vrn-1* or *Vrn-1h* versus the average of the lines carrying the *Vrn-1f* and *Vrn-1g* alleles. Samples were collected when the plants were two weeks old (B) and six weeks old (C).

## Tables


**Table 1.** Survival, relative conductivity and  $F_v/F_m$  in mutant ( $mvp-2/mvp-2 = m/m$ ) and plants carrying at least one wild type *VRN-1* allele ( $Mvp-2/- = M/-$ ). Sixty-day old plants were acclimated at 10 °C for 19 days, 4 °C for 12 days, and -2 °C for 12 h before the freezing treatment. *P* values correspond to factorial ANOVAs including temperature, genotype, and their interactions in the model. Experimental conditions are described in Material and Methods.

Freezing temperatures	-9 °C		-11 °C		Factorial ANOVA
Trait	<i>M/-</i>	<i>m/m</i>	<i>M/-</i>	<i>m/m</i>	<i>P</i> between genotypes
$F_v/F_m$	0.56	0.78	0.56	0.81	<0.0001
Relative conductivity (%)	12.0	10.1	17.3	11.9	0.05
No. of plants that regrew per total (%)	0/50 (0%)	26/30 (87%)	0/59 (0%)	11/24 (46%)	NA*


\* All *Mvp-2/-* plants failed to re-grow and therefore there is no variance within this class to perform an ANOVA. The differences between *mvp-2/mvp-2* and *Mvp-2/-* were obvious.




**Fig. 1:** Quantitative RT-PCR analysis of transcript levels of the *CBF* genes present at the *FR-2* locus relative to the *ACTIN* endogenous control. Samples were collected from leaves of 4-week old *mvp-2/mvp-2* and *Mvp-2/-* plants (20 °C) and again one day later at the same time following an 8 hour cold treatment at 4 °C. Values in the Y axes were normalized and calibrated using the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen 2001). The same calibrator was used for all genes so scales are comparable across genes. Values are averages of 8 biological replications  $\pm$  SE of the means. The inset shows *CBF14* transcript levels which were significantly higher than the other genes at this locus. *P* values for the differences between *mvp/mvp* and *Mvp-/-* after the cold treatment were calculated using ANOVA and are indicated by \*:  $P < 0.05$  or \*\*:  $P < 0.01$ .



**Fig. 2.** Quantitative RT-PCR transcript levels of *COR14b* relative to *TEF1* endogenous control. Plants were 4 weeks old at the beginning of the experiment and were exposed to 4 °C for 12days. Values in the Y axes are normalized and calibrated values using the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen 2001). Homozygous *mvp-2/mvp-2* plants (null *VRN-1*) are indicated by black squares and lines and *Mvp-1/-* plants (1 or 2 *VRN-1* copies) by gray triangles and lines. Values are averages of 8 biological replications in the untransformed scale  $\pm$  SE of the means. *P* values were calculated using ANOVA of  $\log(n+1)$  transformed values for each time point (\*\*:  $P < 0.01$ ).



**Fig. 3.** Time courses of *VRN-1*, and *COR* genes *COR14b*, *DHN5*, and *DHN8* transcripts in *mvp-2/mvp-2* plants homozygous for a *VRN-1* deletion (*m*) and *Mvp-2/-* plants with one or two functional *VRN-1* copies (*M*). The presence of a faint *VRN-1* hybridization in the mutant sample from 87 days is suspected to be due to cross contamination.



**Fig. 4.** *VRN-1* and *COR14b* transcript levels in a set of *T. monococcum* lines differing in *VRN-1* expression under short days. Abbreviations: W= warm conditions, C= decrease from 18 °C to 6 °C (occurring at daybreak), 2W= two-week old, 6W= six-week old, SD = short day, LD = long day, wt= wild type, *Vrn-1f*= allele with 1-bp deletion in the CArG box coupled with *VRN-1* intron 1 insertion, *Vrn-1g* = allele with 34-bp deletion encompassing the CArG box. *Vrn-1h* = allele with insertion in *VRN-1* intron 1. The three accessions with the *Vrn-1f* allele carry the dominant *Vrn-2* allele whereas all the other accessions carry non-functional *vrn-2* alleles. All the accessions have a spring growth habit. **A**) mRNA blot analyses of three genotypes per promoter class (indicated by different PI numbers). Arrows and arrowheads identify the presence and absence, respectively, of *COR14b* transcripts in 6W plants in the *Vrn-1f* and *Vrn-1g* natural mutants under SD and LD. **B-C**) qRT-PCR validation of *VRN-1* and *COR14b* transcript levels relative to *ACTIN* endogenous control at 4 °C. Values in the Y axes are normalized and calibrated values using the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen 2001). Lines carrying the wild type allele (*vrn-1*) or the *Vrn-1h* (spring, not induced at SD) are indicated by black bars and lines carrying the *Vrn-1f* and *Vrn-1g* alleles (spring, induced at SD) by gray bars. Values are averages of 5 biological replications  $\pm$  SE of the means. *P* values were calculated using contrasts between either *vrn-1* or *Vrn-1h* versus the average of the lines carrying the *Vrn-1f* and *Vrn-1g* alleles. Samples were collected when the plants were two weeks old (**B**) and six weeks old (**C**).